arXiv:1612.02644v1 [gr-qc] 8 Dec 2016

Exact solution of scalar field in Schwarzschild spacetime: bound state and scattering state

Wen-Du Li,[a,b] **Yu-Zhu Chen,**[b] **and Wu-Sheng Dai**[b,c,*]

[a] *Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin, 300071, P. R. China*

[b] *Department of Physics, Tianjin University, Tianjin 300072, P.R. China*

[c] *LiuHui Center for Applied Mathematics, Nankai University & Tianjin University, Tianjin 300072, P.R. China*

Abstract: Exact solutions of a scalar field in the Schwarzschild spacetime are presented. For bound states, we obtain the exact bound-state wave function and eigenvalue; for scattering states, we obtain the exact scattering wave function and phase shift. By virtue of the exact solutions, we give a direct calculation for the discontinuous jump on horizon for massive scalar fields, while in literature such a jump is obtained by an asymptotic solution by an analytic extension technique for massless cases. Besides the exact solution, we also provide an asymptotic expression for bound-state eigenvalues. Additionally, as an alternative approach, we develop an integral equation method for the calculation of scattering wave functions and scattering phase shifts.

*daiwusheng@tju.edu.cn.

Contents

1 Introduction **1**

2 Scalar field in Schwarzschild spacetime **3**
2.1 Field equation 3
2.2 Boundary condition 3

3 Converting Scalar field equation to confluent Heun equation **4**
3.1 Confluent Heun equation 4
3.2 Boundary condition 5

4 Bound state **5**
4.1 Bound-state wave function 5
4.2 Bound-state eigenvalue 7

5 Scattering state: Exact solution of wave function and phase shift **8**
5.1 Scattering wave function: Regular solution 8
5.1.1 Exact solution of regular solution 8
5.2 Scattering wave function: Irregular solution 9
5.2.1 Scattering boundary condition: Asymptotic behavior 9
5.2.2 Exact solution of outgoing wave function: Outgoing Eddington-Finkelstein coordinate 9
5.2.3 Exact solution of ingoing wave function: Ingoing Eddington-Finkelstein coordinate 11
5.3 Scattering phase shift 12
5.4 Asymptotic behavior of scattering wave function 12

6 Bound-state eigenvalue: Exact solution **12**

7 Discontinuous jump on horizon: Jump condition **13**
7.1 Jump condition: Outgoing wave function 13
7.2 Jump condition: Ingoing wave function 13

8 Discontinuous jump on horizon: Comparison with Hawking and Damour-Ruffini methods **14**
8.1 Hawking method: Brief review 14
8.2 Damour-Ruffini method: Brief review 15
8.3 Direct calculation through exact solution: Comparison 16

9 Integral equation method: Alternative approach for scattering wave function and phase shift **16**
9.1 Integral equation 16

9.1.1 Outer horizon 17
9.1.2 Wave function near horizon: Outer horizon 18
9.1.3 Inner horizon 19
9.1.4 Wave function near horizon: Inner horizon 20
9.2 Connect wave functions by jump condition on horizon 20
9.2.1 Jump condition 20
9.2.2 Relative scattering amplitude 20
9.3 Alternative expression of scattering phase shift 21
9.3.1 Scattering phase shift 22
9.3.2 Scattering phase shift: zeroth-order and first-order 23

10 Conclusion 24

1 Introduction

In this paper, we provide an exact solution of a scalar field in the Schwarzschild spacetime, including scattering and bound-state solutions.

A massive scalar field Φ with mass μ in the background of the Schwarzschild spacetime,

$$ds^2 = -\left(1-\frac{2M}{r}\right)dt^2 + \left(1-\frac{2M}{r}\right)^{-1}dr^2 + r^2 d\theta^2 + r^2\sin^2\theta d\phi^2, \tag{1.1}$$

is described by the scalar equation [1]

$$\left(\frac{1}{\sqrt{-g}}\frac{\partial}{\partial x^\mu}\sqrt{-g}g^{\mu\nu}\frac{\partial}{\partial x^\nu} - \mu^2\right)\Phi = 0. \tag{1.2}$$

The key problem of solving the equation is that there are two singular points in the background spacetime: $r = 0$ and $r = 2M$ [2]. At these singular points the differential equation (1.2) is undefined and needs two boundary conditions put in by hand respectively. Nevertheless, besides these two boundary conditions on the two singular points, there is also a boundary condition at $r \to \infty$ — the natural boundary of the spacetime. Therefore, the difficulty we encountered is that Eq. (1.2) is a second order differential equation, the solution must satisfy these three boundary conditions simultaneously. In this paper, we show how to overcome this difficulty.

On the horizon, there is a discontinuous jump of wave function. The calculation of the Hawking radiation relies on the magnitude of the jump. In order to calculate the jump, Hawking [3] and Damour and Ruffini [4], starting from an asymptotic wave function, introduce an analytic extension technique.

Instead of the analytic extension treatment, in this paper, starting from an exact wave function rather than an approximate asymptotic one, we calculate the discontinuous jump directly.

The exact solutions of the bound-state wave function, the bound-state eigenvalue, scattering wave function, the scattering phase shift, and the exact jump of wave function on the horizon are obtained in the present paper.

Besides the exact solutions, we also provide some approximate results. An asymptotic expression for the eigenvalue of bound states is presented. Moreover, we establish an integral equation method for the calculation of the scattering wave function and the phase shift. Using the integral equation method, we calculate the first two orders contributions of the scattering wave function and the scattering phase shift. The integral equation method constructed in the present paper is a systematic method for solving the problem of scattering by a spacetime and can be applied to other scattering problems in gravity.

In black hole theory, the study of scattering plays an important role [1, 5]. There are many studies on scattering, such as the asymptotic tail [6] and complex angular momenta of scalar scattering [7]. The absorption cross section of regular black holes which have event horizons but not singularities is discussed [8]. Many approximate methods are developed, such as the Born approximation of massless scalar scattering [9], phase-integral method for scalar scattering [10], the propagation of a massive vector field [11], massive Dirac field scattering [12], the absorption cross section for scalar scattering [13], orbiting scattering of massless spin 0, 1 and 2 particles [14], massive spin-half scattering [15], the WKB approximation for massive Dirac field scattering [16] in the Schwarzschild spacetime, a massive scalar scattering in the Reissner-Nordström spacetime [17], massless planar scalar waves scattered by a charged nonrotating black hole [18], scattering by a deformed non-rotating black hole [19], and massless scalar scattering by a Kerr black hole [20]. Scattering of spin fields and vector fields in curved spacetime is also studied: the analogue of the Mott formula for scattering in a Coulomb background and in the Dirac scattering by a black hole [21], massive spin-2 fluctuations of Schwarzschild and slowly rotating Kerr black holes [22], the internal stationary state of a black hole for massless Dirac fields [23], the quasinormal modes of electromagnetic and gravitational perturbations of a Schwarzschild black hole in an asymptotically anti-de Sitter spacetime [24], and the quasinormal mode frequencies for the massless Dirac field in Schwarzschild-AdS spacetime [25]. Scattering on arbitrary dimensional black holes and on black holes with a cosmological constant is considered [26]. Scattering between two black holes is numerically studied [26]. The scalar field perturbations of the $4+1$-dimensional Schwarzschild black hole immersed in a Gödel universe by the Gimon-Hashimoto solution is described [27]. Scattering method can be used in the calculation of the Hawking radiation. A systematic scattering method for the Hawking radiation is developed by Damour and Ruffini [4, 28]. The Hawking radiation of a Reissner–Nordström–de Sitter black hole [29], the scalar particle Hawking radiation of a BTZ black hole [30], the charged Dirac particle Hawking radiation of the Kerr-Newman black hole [31], and the distribution for particles emitted by a black hole [32] are discussed by the Damour–Ruffini method. The Dirac particle Hawking radiation of the Kerr black hole [33] and of the BTZ black hole [34] are calculated by the WKB approximation. The Hawking radiation of acoustic black holes is discussed [35]. The renormalized expectation values $\langle T_{ab} \rangle$ of the relevant energy-momentum tensor operator of a massless scalar field in the Schwarzschild spacetime is calculated [36]. Some exact solutions are also obtained. The analytical solution of the

Regge-Wheeler equation and the Teukolsky radial equation is obtained in Ref. [37]. In Ref. [38], using the truncation condition of the confluent Heun function, the authors calculate resonant frequencies for a charged scalar field in a dyonic black hole background, and the asymptotic form of the scattering wave function and Hawking radiation are presented. An exact solution of the Klein–Gordon equation in Kerr–Newman spacetime is calculated, but in which the scattering and bound-state boundary condition are not taken into account [39].

In section 2, the equation of a scalar field in a Schwarzschild spacetime and the corresponding boundary conditions for bound states and scattering states are given. In section 3, as a key step, we convert the scalar field equation in the Schwarzschild spacetime into a confluent Heun equation. In section 4, we provide an exact solution of the bound-state wave function and an explicit asymptotic expression for the bound-state eigenvalue; the exact solution of the bound-state eigenvalue will be given in section 6. In section 5, an exact solution of the scattering wave function and an exact solution of the scattering phase shift are given. In section 6, we give an exact solution of the bound-state eigenvalue. In section 7, based on the exact wave functions given in the above sections, we determine the jump condition of the ingoing and outgoing wave functions on the horizon. In section 8, we compare our result of the discontinuous jump on the horizon with the Hawking and Damour-Ruffini treatments. In section 9, we develop an alternative approach for the calculation of the scattering wave function and the phase shift, which is an integral equation method. The conclusions are summarized in section 10.

2 Scalar field in Schwarzschild spacetime

2.1 Field equation

The Schwarzschild spacetime is spherically symmetric, so we can perform a partial-wave expansion [1]:

$$\Phi\left(x^{\mu}\right)=\sum_{l=0}^{\infty}\left(2l+1\right)e^{-i\omega t}P_{l}\left(\cos\theta\right)\phi_{l}\left(r\right), \tag{2.1}$$

where $\phi_{l}\left(r\right)$ is the radial wave function satisfying the radial equation

$$\left[\frac{1}{r^{2}}\left(1-\frac{2M}{r}\right)\frac{d}{dr}r^{2}\left(1-\frac{2M}{r}\right)\frac{d}{dr}+\omega^{2}-\left(1-\frac{2M}{r}\right)\mu^{2}-\left(1-\frac{2M}{r}\right)\frac{l\left(l+1\right)}{r^{2}}\right]\phi_{l}\left(r\right)=0. \tag{2.2}$$

There exist three singularities, $r=0$, $2M$, and ∞, in the radial equation. $r=2M$ is the horizon of the Schwarzschild spacetime and $r\rightarrow\infty$ is the natural boundary of space.

2.2 Boundary condition

The first problem we encountered is how to impose three boundary conditions on one second order differential equation at a time.

At singular points, one needs to impose boundary conditions by hand, since the differential equation is undefined at these singular points.

The equation of a scalar field in the Schwarzschild spacetime, Eq. (2.2), besides the natural boundary at ∞, has two singularities, $r = 0$ and $r = 2M$, rather than that of the common singular potential in quantum mechanics which often have only one single singularity at $r = 0$.

The first problem encountered is that if there are n singular points and a natural boundary at ∞, one needs to impose $n + 1$ boundary conditions: n for singular points and one for ∞. Nevertheless, the dynamical equation (2.2) is a second order differential equation, so, at any one time one can at most impose two boundary conditions. Once the number of the singular points is more than 1, we must insure that there is no contradiction.

In our problem, we have to impose three boundary conditions at $r = 0$, $r = 2M$, and $r \to \infty$, respectively. The radial equation (2.2) is a second order differential equation, so only two boundary conditions can be imposed at a time. In order to impose these three boundary conditions on one second order differential equation, we will, respectively, solve the equation with the boundary conditions at $r = 0$ and $r = 2M$ and solve the equation with the boundary conditions at $r = 2M$ and $r \to \infty$. Finally, we connect these two solutions together properly with an appropriate connection condition.

The boundary conditions are as follows.

The boundary condition at the two singular points of the Schwarzschild spacetime, $r = 0$ and $r = 2M$, are

$$|\phi_l(0)| < \infty, \tag{2.3}$$

$$|\phi_l(2M)| < \infty, \tag{2.4}$$

i.e., $\phi_l(r)$ must be finite at the singular points.

The boundary condition at $r \to \infty$ determines the solution whether a bound state or a scattering state:

$$\phi_l(r)|_{r\to\infty} = 0, \qquad \text{bound state}, \tag{2.5}$$

$$\phi_l(r)|_{r\to\infty} = \phi_l^{\infty}(r), \quad \text{scattering state}, \tag{2.6}$$

where $\phi_l^{\infty}(r)$ is the large-distance asymptotics of the solution of the radial equation.

Scattering by a Schwarzschild spacetime is essentially a kind of long-range scattering [5]. Recall that for potential scattering, the large-distance asymptotics of the solution of the radial equation, $\phi_l^{\infty}(r)$, are the same for all short-range potential scattering, but for long-range potential scattering [40, 41], like that in our case, $\phi_l^{\infty}(r)$ is determined by the potential and different potentials have different asymptotic solutions [42, 43].

3 Converting Scalar field equation to confluent Heun equation

3.1 Confluent Heun equation

The key step in solving the scalar field equation in the Schwarzschild spacetime is to convert the radial equation (2.2) into a confluent Heun equation, also called the generalized spheroidal equation [44].

By the variable substitution $z = r/M - 1$, the radial equation (2.2) can be converted into a confluent Heun equation

$$\frac{d}{dz}\left(z^2-1\right)\frac{d}{dz}y\left(z\right)+\left[-p^2\left(z^2-1\right)+2p\beta z-\lambda-\frac{m^2+s^2+2msz}{z^2-1}\right]y\left(z\right)=0, \tag{3.1}$$

where the parameters

$$m = s = i2M\sqrt{\eta^2+\mu^2}, \tag{3.2}$$

$$\beta = -i2M\eta - iM\frac{\mu^2}{\eta}, \tag{3.3}$$

$$p = iM\eta, \tag{3.4}$$

$$\lambda = l\left(l+1\right) - 8\eta^2M^2 - 6\mu^2M^2 \tag{3.5}$$

with $\eta = \sqrt{\omega^2-\mu^2}$.

The confluent Heun equation (3.1) has three singular points: $z = -1,\, 1,\, \infty$ [44]. These three singular points just correspond to the three singular points of the Schwarzschild spacetime, $r = 0,\, 2M,\, \infty$.

3.2 Boundary condition

The boundary condition of $\phi_l\left(r\right)$ is then converted into a boundary condition of $y\left(z\right)$.

The boundary conditions at two singular points $r = 0$ and $r = 2M$, Eqs. (2.3) and (2.4), become

$$\left|y\left(-1\right)\right| < \infty, \tag{3.6}$$

$$\left|y\left(1\right)\right| < \infty. \tag{3.7}$$

The boundary conditions for bound states and scattering states at $r \to \infty$ then become

$$y\left(z\right)|_{z\to\infty} = 0, \qquad \text{bound state}, \tag{3.8}$$

$$y\left(z\right)|_{z\to\infty} = y^{\infty}\left(z\right), \quad \text{scattering state}, \tag{3.9}$$

where $y^{\infty}\left(z\right)$ is the large-distance asymptotics of the solution of the confluent Heun equation Eq. (3.1).

4 Bound state

4.1 Bound-state wave function

For bound states, let

$$\eta = ik, \tag{4.1}$$

so that for bound states $\eta = -k^2 < 0$ with k a real number. Then

$$m = s = -2M\sqrt{k^2 - \mu^2}, \tag{4.2}$$

$$\beta = \frac{M}{k}\left(2k^2 - \mu^2\right), \tag{4.3}$$

$$p = -kM, \tag{4.4}$$

$$\lambda = l\left(l+1\right) + 2M^2\mu^2 + 8M^2\left(k^2 - \mu^2\right). \tag{4.5}$$

The bound-state boundary condition is $\phi_l\left(\infty\right) = 0$, or, $y\left(\infty\right) = 0$, given by Eqs. (2.5) and (3.8).

Next, we solve the radial equation (2.2) with the boundary conditions at $r = 0$ and $r = 2M$ ($z = -1$ and $z = 1$) and the boundary conditions at $r = 2M$ and $r \to \infty$ ($z = 1$ and $z \to \infty$), respectively.

At $r = 0$ and $r = 2M$ ($z = -1$ and $z = 1$) the radial equation (2.2) with the boundary conditions (2.3) and (2.4), corresponding to the confluent Heun equation (3.1) with the boundary conditions (3.6) and (3.7), yields a solution $\Xi\left(p, \beta, z\right)$ satisfying $\left|\Xi\left(p, \beta, -1\right)\right| < \infty$ and $\left|\Xi\left(p, \beta, 1\right)\right| < \infty$, called the angular generalized spheroidal function of p-type (AGSF) [44].

At $r = 2M$ and $r \to \infty$ ($z = 1$ and $z = \infty$) the radial equation (2.2) with the boundary conditions (2.4) and (2.5), corresponding to the confluent Heun equation (3.1) with the boundary conditions (3.7) and (3.8), yields another solution $\Pi\left(p, \beta, z\right)$ satisfying $\left|\Pi\left(p, \beta, 1\right)\right| < \infty$ and $\left|\Pi\left(p, \beta, \infty\right)\right| = 0$, called the radial generalized spheroidal function of p-type (RGSF) [44].

The solution, however, must simultaneously satisfy all the three boundary conditions at $r = 0$ ($z = -1$), $r = 2M$ ($z = 1$), and $r \to \infty$ ($z \to \infty$). This requires the connection condition

$$\Xi\left(p, \beta, z\right) = \Pi\left(p, \beta, z\right). \tag{4.6}$$

The bound-state wave solution which satisfies the connection condition (4.6) and the bound-state boundary conditions (2.3), (2.4), and (2.5) is [44]

$$\begin{aligned}\Xi\left(p, \beta, z\right) = {} & N\left(z-1\right)^{(m+s)/2}\left(z+1\right)^{(m-s)/2} e^{-p(1+z)} \\ & \times \mathrm{Hc}^{\,(a)}\left(p, -\beta + m + 1, m + s + 1, m - s + 1, \sigma; \frac{z+1}{2}\right),\end{aligned} \tag{4.7}$$

where $\mathrm{Hc}^{\,(a)}\left(p, \alpha, \gamma, \delta, \sigma; z\right)$ is the angular confluent Heun function, N is the normalization constant, and $\sigma = \lambda + 2p\left(-2\beta + m + s + 1\right) - m\left(m+1\right)$ with the restriction

$$m + s \geq 0,\ \ m - s \geq 0,\ \ p \geq 0,\ \ \beta \in R. \tag{4.8}$$

The restrictions (4.4) and (4.8) require that

$$k < 0. \tag{4.9}$$

The bound-state wave function can be then expressed as

$$\begin{aligned}\Xi(k,r) &= N\left(\frac{r}{M}-2\right)^{-2M\sqrt{k^2-\mu^2}}e^{kr}\\ &\times \mathrm{Hc}^{(a)}\left(-Mk,1+\frac{M}{k}\left(\mu^2-2k^2\right)-2M\sqrt{k^2-\mu^2},1-4M\sqrt{k^2-\mu^2},1,\sigma;\frac{r}{2M}\right)\\ &= NM^{2M\sqrt{k^2-\mu^2}}e^{kr-2M\sqrt{k^2-\mu^2}\ln(r-2M)}\\ &\times \mathrm{Hc}^{(a)}\left(-Mk,1-M\left(k+\sqrt{k^2-\mu^2}\right)^2,1-4M\sqrt{k^2-\mu^2},1,\sigma;\frac{r}{2M}\right),\end{aligned} \tag{4.10}$$

where N is the normalization constant and

$$\sigma = l(l+1)+2M^2\left(k^2+\mu^2\right)+8M^2\left(\sqrt{k^2-\mu^2}+\frac{k}{2}\right)^2-2M\left(k-\sqrt{k^2-\mu^2}\right). \tag{4.11}$$

An explicit asymptotic expression of the eigenvalue $-k^2$ will be given later in section 4.2); an implicit exact expression of the eigenvalue $-k^2$ is given by Eq. (6.2) in section 6.

4.2 Bound-state eigenvalue

To implement the requirement (4.6), we must take the value of the parameter λ in Eq. (3.1) as discrete values [44]:

$$\lambda = \lambda_n(p,\beta) \tag{4.12}$$

with n an integer. Together with Eqs. (4.3), (4.4), and (4.5), we have

$$l(l+1)+2M^2\mu^2+8M^2\left(k^2-\mu^2\right) = \lambda_n\left(-Mk,\frac{M}{k}\left(2k^2-\mu^2\right)\right). \tag{4.13}$$

The energy eigenvalue $-k^2$ can be solved from Eq. (4.13).

Before solving the exact eigenvalue which will be given by analyzing the analytic property of the S-Matrix of scattering in section 6, we now give an asymptotic expression of the eigenvalue of bound states.

For a large $p=-Mk$, the eigenvalue λ_n has the following asymptotics [44]:

$$\begin{aligned}\lambda_n &= -2Mk\left[2\chi-\frac{M}{k}\left(2k^2-\mu^2\right)\right]+\left\{-2\chi\left[\chi-\frac{M}{k}\left(2k^2-\mu^2\right)\right]+\frac{1}{2}\left[8M^2\left(k^2-\mu^2\right)-1\right]\right\}\\ &+\frac{1}{2Mk}\left\{\chi\left[\chi-\frac{M}{k}\left(2k^2-\mu^2\right)\right]^2-\left(\chi^2-\frac{1}{4}\right)\left[\chi-\frac{M}{k}\left(2k^2-\mu^2\right)\right]+\frac{1}{4}\chi\left[1+8M^2\left(k^2-\mu^2\right)\right]\right\}\\ &+O\left(\frac{1}{(Mk)^2}\right),\end{aligned} \tag{4.14}$$

where

$$\chi = n+\frac{1}{2} \tag{4.15}$$

with n an integer. The eigenvalue $\eta^2=-k^2$ can be solved from Eqs. (4.14) and (4.13) directly.

For instance, up to the first order of $\frac{1}{Mk}$ for simplicity, by Eqs. (4.13), (4.14), and (4.15), we can obtain the eigenvalue. Solving

$$\begin{aligned} & l\left(l+1\right)+2M^{2}\mu^{2}+8M^{2}\left(k^{2}-\mu^{2}\right) \\ & =-2Mk\left[2\left(n+\frac{1}{2}\right)-\frac{M}{k}\left(2k^{2}-\mu^{2}\right)\right]+\left\{-2\left(n+\frac{1}{2}\right)\left(n+\frac{1}{2}-\frac{M}{k}\left(2k^{2}-\mu^{2}\right)\right)\right. \\ & \left.+\frac{1}{2}\left[8M^{2}\left(k^{2}-\mu^{2}\right)-1\right]\right\} \end{aligned} \tag{4.16}$$

gives

$$k=-\frac{2n+1}{2n\left(n+1\right)+l\left(l+1\right)+1}\mu^{2}M^{2}. \tag{4.17}$$

Then the eigenvalue reads

$$\eta^{2}=-k^{2}=-\left[\frac{2n+1}{2n\left(n+1\right)+l\left(l+1\right)+1}\right]^{2}\mu^{4}M^{4}. \tag{4.18}$$

An exact implicit expression of the eigenvalue of bound states will be given by analyzing the analytic property of the S-Matrix of scattering in section 6.

5 Scattering state: Exact solution of wave function and phase shift

For scattering states,

$$p=ic,\quad \beta=-i\zeta, \tag{5.1}$$

where $c=M\eta$ and $\zeta=2M\eta+M\mu^{2}/\eta$.

For scattering states, similarly to bound states, we also solve the radial equation (2.2) with the boundary conditions at $r=0$ and $r=2M$ and the boundary conditions at $r=2M$ and $r\to\infty$, respectively, replacing the bound-state boundary condition (2.5) with the scattering-state boundary condition (2.6) at $r\to\infty$.

5.1 Scattering wave function: Regular solution

For scattering in the Schwarzschild spacetime, the regular solution is a solution satisfying the boundary condition at $r=0$ and $r=2M$. As a comparison, in common central potential scattering the regular solution usually only needs to satisfy the boundary condition at one singular point $r=0$ [42].

5.1.1 Exact solution of regular solution

At $r=0$ and $r=2M$, the solution of the radial equation (2.2) still needs to satisfy the boundary conditions (2.3) and (2.4), but the parameters p and β given by Eq. (5.1) are now pure imaginary numbers. The solution now is the angular generalized spheroidal function of c-type, $\Xi\left(c,\zeta,z\right)$, which can be achieved by substituting $\eta=ik$ into Eq. (4.10) [44]:

$$\begin{aligned} \Xi\left(\eta,r\right) & =NM^{-i2M\sqrt{\eta^{2}+\mu^{2}}}e^{-i\left[\eta r-2M\sqrt{\eta^{2}+\mu^{2}}\ln(r-2M)\right]} \\ & \times\mathrm{Hc}^{(a)}\left(iM\eta,1+i\frac{M}{\eta}\left(\eta+\sqrt{\eta^{2}+\mu^{2}}\right)^{2},1+i4M\sqrt{\eta^{2}+\mu^{2}},1,\sigma;\frac{r}{2M}\right), \end{aligned} \tag{5.2}$$

where

$$\sigma = l\left(l+1\right) - 2M^2\left(\eta^2-\mu^2\right) - 8M^2\left(\frac{\eta}{2}+\sqrt{\eta^2+\mu^2}\right)^2 + i2M\left(\eta-\sqrt{\eta^2+\mu^2}\right). \quad (5.3)$$

This solution is an analogue of the regular solution in quantum-mechanical scattering theory. The regular solution in quantum-mechanical central potential scattering satisfies the boundary condition at only *one* singular point $r=0$ [45]. In our problem, the regular solution (5.2) simultaneously satisfies two boundary conditions at two singular points $r=0$ and $r=2M$.

5.2 Scattering wave function: Irregular solution

For scattering in the Schwarzschild spacetime, the irregular solution is the solution satisfying the boundary conditions at $r=2M$ and $r\to\infty$. As a comparison, in quantum-mechanical central potential scattering the irregular solution usually only needs to satisfy the boundary condition at $r\to\infty$ [42].

At $r=2M$ and $r\to\infty$, the solution of the radial equation (2.2) still needs to satisfy the boundary condition (2.4), but the bound-state boundary condition (2.5) is replaced by the scattering boundary condition (2.6) and the parameters p and β are taken as imaginary numbers.

5.2.1 Scattering boundary condition: Asymptotic behavior

Before seeking exact solutions, we first investigate the large-distance asymptotic behavior of the scattering wave function. The asymptotic solution of the scattering wave function will serve as boundary conditions for scattering states [40–42].

The solution of Eq. (2.2) satisfying the scattering boundary condition at $r\to\infty$ is the radial generalized spheroidal function (RGSF) of c-type, $\Pi\left(\eta,r\right)$ [44].

The large-distance asymptotics of the RGSF of c-type is [44]

$$\Pi^{(1)}\left(\eta,r\right) \overset{r\to\infty}{\sim} \frac{1}{r}\exp\left(i\left[\eta\left(r-M\right)+2\eta M\ln\frac{r}{M}+M\frac{\mu^2}{\eta}\ln\left(\frac{r}{M}-1\right)+\chi\left(\eta,M\right)\right]\right), \quad (5.4)$$

$$\Pi^{(2)}\left(\eta,r\right) \overset{r\to\infty}{\sim} \frac{1}{r}\exp\left(-i\left[\eta\left(r-M\right)+2\eta M\ln\frac{r}{M}+M\frac{\mu^2}{\eta}\ln\left(\frac{r}{M}-1\right)+\chi\left(\eta,M\right)\right]\right), \quad (5.5)$$

where $\chi\left(\eta,M\right)$ is the phase of the RGSF of c-type [44]. For high-energy scattering, we have [44]

$$\chi\left(\eta,M\right) = -2M\eta\ln 2 - \frac{l\pi}{2} - 2M\eta\pi + O\left(\frac{1}{M\eta}\right). \quad (5.6)$$

5.2.2 Exact solution of outgoing wave function: Outgoing Eddington-Finkelstein coordinate

In order to solve the scattering wave function, we use the Eddington-Finkelstein coordinate.

The scalar equation with outgoing Eddington-Finkelstein coordinate. To solve the outgoing wave function, we use the outgoing Eddington-Finkelstein coordinate (u, r) with $u = t - r_*$ [46]. By the outgoing Eddington-Finkelstein coordinate, we have

$$ds^2 = -\left(1 - \frac{2M}{r}\right) du^2 - 2dudr + r^2 d\theta^2 + r^2 \sin^2\theta d\phi^2. \tag{5.7}$$

The scalar equation (1.2) then becomes

$$\left[-\frac{1}{r^2}\frac{\partial}{\partial u}r^2\frac{\partial}{\partial r} - \frac{1}{r^2}\frac{\partial}{\partial r}r^2\frac{\partial}{\partial u} + \frac{1}{r^2}\frac{\partial}{\partial r}r^2\left(1 - \frac{2M}{r}\right)\frac{\partial}{\partial r} + \frac{L^2}{r^2} - \mu^2\right]\Phi = 0. \tag{5.8}$$

By means of the variable separation

$$\Phi = \beta(u, r) Y_{lm}(\theta, \phi), \tag{5.9}$$

we arrive at

$$\left[-\frac{1}{r^2}\frac{\partial}{\partial u}r^2\frac{\partial}{\partial r} - \frac{1}{r^2}\frac{\partial}{\partial r}r^2\frac{\partial}{\partial u} + \frac{1}{r^2}\frac{\partial}{\partial r}r^2\left(1 - \frac{2M}{r}\right)\frac{\partial}{\partial r} - \frac{l(l+1)}{r^2} - \mu^2\right]\beta(u, r) = 0. \tag{5.10}$$

This equation can be solved exactly.

Outer horizon. Outside the horizon, the outgoing solution of Eq. (5.10) reads [44]

$$\begin{aligned}\beta^{outer}(u, r) = {} & e^{-i\sqrt{\eta^2+\mu^2}u} e^{-i\sqrt{\eta^2+\mu^2}r} e^{i\eta r} e^{i2M\eta(1-r/M)} \\ & \times \mathrm{Hc}^{(a)}\left(-iM\eta, i2M\left(\eta + \sqrt{\eta^2+\mu^2}\right) + iM\frac{\mu^2}{\eta} + 1, 1, 1 + i4M\sqrt{\eta^2+\mu^2}, \sigma^{outer}, 1 - \frac{r}{2M}\right),\end{aligned} \tag{5.11}$$

where $\mathrm{Hc}^{(a)}(\alpha, \beta, \gamma, \delta, \sigma, z)$ is the confluent Heun function and

$$\sigma^{outer} = l(l+1) + 4M^2\left(\eta^2 + \frac{\mu^2}{2}\right) - i2M\left(\eta + \sqrt{\eta^2+\mu^2}\right). \tag{5.12}$$

Inner horizon. Inside the horizon, the outgoing solution of Eq. (5.10) reads [44]

$$\begin{aligned}\beta^{inner}(u, r) = {} & e^{-i\sqrt{\eta^2+\mu^2}u} e^{-i\sqrt{\eta^2+\mu^2}r} e^{i\eta r} (r - 2M)^{-i4\sqrt{\eta^2+\mu^2}M} e^{i2M\eta(1-r/M)} \\ & \times \mathrm{Hc}^{(a)}\left(-iM\eta, i2M\left(\eta - \sqrt{\eta^2+\mu^2}\right) + iM\frac{\mu^2}{\eta} + 1, 1, 1 - i4M\sqrt{\eta^2+\mu^2}, \sigma^{inner}, 1 - \frac{r}{2M}\right),\end{aligned} \tag{5.13}$$

where

$$\sigma^{inner} = l(l+1) + 4M^2\left(\eta^2 + \frac{\mu^2}{2}\right) - i2M\left(\eta - \sqrt{\eta^2+\mu^2}\right). \tag{5.14}$$

Scattering wave function. For scattering, we only concern ourselves with the wave function outside the horizon, so the exact irregular solution reads

$$\Pi^{(1)}(\eta, r) = \beta^{outer}(u, r). \tag{5.15}$$

5.2.3 Exact solution of ingoing wave function: Ingoing Eddington-Finkelstein coordinate

The scalar equation with ingoing Eddington-Finkelstein coordinate. For the ingoing wave function, instead of the outgoing Eddington-Finkelstein coordinate, we use the ingoing Eddington-Finkelstein coordinate (v,r) with $v=t+r_*$ [46]. By the ingoing Eddington-Finkelstein coordinate, we have

$$ds^2=-\left(1-\frac{2M}{r}\right)dv^2+2dvdr+r^2d\theta^2+r^2\sin^2\theta d\phi^2. \tag{5.16}$$

Eq. (1.2) then becomes

$$\left[\frac{1}{r^2}\frac{\partial}{\partial v}r^2\frac{\partial}{\partial r}+\frac{1}{r^2}\frac{\partial}{\partial r}r^2\frac{\partial}{\partial v}+\frac{1}{r^2}\frac{\partial}{\partial r}r^2\left(1-\frac{2M}{r}\right)\frac{\partial}{\partial r}-\frac{L^2}{r^2}-\mu^2\right]\Phi=0 \tag{5.17}$$

By means of the variable separation

$$\Phi=\alpha\left(v,r\right)Y_{lm}\left(\theta,\phi\right), \tag{5.18}$$

we arrive at

$$\left[\frac{1}{r^2}\frac{\partial}{\partial v}r^2\frac{\partial}{\partial r}+\frac{1}{r^2}\frac{\partial}{\partial r}r^2\frac{\partial}{\partial v}+\frac{1}{r^2}\frac{\partial}{\partial r}r^2\left(1-\frac{2M}{r}\right)\frac{\partial}{\partial r}-\frac{l\left(l+1\right)}{r^2}-\mu^2\right]\alpha\left(v,r\right)=0. \tag{5.19}$$

Outer horizon. Outside the horizon, the ingoing solution of Eq. (5.19) reads [44]

$$\begin{aligned}\alpha^{outer}\left(v,r\right)=&e^{-i\sqrt{\eta^2+\mu^2}v}e^{i\sqrt{\eta^2+\mu^2}r}e^{i\eta r}\left(r-2M\right)^{i4\sqrt{\eta^2+\mu^2}M}e^{i2M\eta(1-r/M)}\\&\times\mathrm{Hc}^{(a)}\left(-iM\eta,i2M\left(\eta+\sqrt{\eta^2+\mu^2}\right)+iM\frac{\mu^2}{\eta}+1,1,1+i4M\sqrt{\eta^2+\mu^2},\sigma^{outer},1-\frac{r}{2M}\right),\end{aligned} \tag{5.20}$$

where σ^{outer} is given by Eq. (5.12).

Inner horizon. Inside the horizon, the ingoing solution of Eq. (5.19) reads [44]

$$\begin{aligned}\alpha^{inner}\left(v,r\right)=&e^{-i\sqrt{\eta^2+\mu^2}v}e^{i\sqrt{\eta^2+\mu^2}r}e^{i\eta r}e^{i2M\eta\left(1-\frac{r}{M}\right)}\\&\times\mathrm{Hc}^{(a)}\left(-iM\eta,i2M\left(\eta-\sqrt{\eta^2+\mu^2}\right)+iM\frac{\mu^2}{\eta}+1,1,1-i4M\sqrt{\eta^2+\mu^2},\sigma^{inner},1-\frac{r}{2M}\right),\end{aligned} \tag{5.21}$$

where σ^{inner} is given by Eq. (5.14).

Scattering wave function. For scattering, we only concern ourselves with the wave function outside the horizon, so the exact irregular solution reads

$$\Pi^{(2)}\left(\eta,r\right)=\alpha^{outer}\left(v,r\right). \tag{5.22}$$

5.3 Scattering phase shift

To calculate the scattering phase shift, we first express the regular solution $\Xi(\eta, r)$ which satisfies the boundary condition at $r = 0$ and $r = 2M$ as a linear combination of the irregular solutions $\Pi^{(1)}(\eta, r)$ and $\Pi^{(2)}(\eta, r)$ which satisfy the boundary condition at $r = 2M$ and $r \to \infty$:

$$\Xi(\eta, r) = C_l i^{l+1} \Pi^{(2)}(\eta, r) + D_l (-i)^{l+1} \Pi^{(1)}(\eta, r). \tag{5.23}$$

Here, the coefficients C_l and D_l is determined by [47]

$$C_l = (-i)^{l+1} \frac{W_r\left[\Pi^{(1)}(\eta, r), \Xi(\eta, r)\right]}{W_r\left[\Pi^{(1)}(\eta, r), \Pi^{(2)}(\eta, r)\right]}, \tag{5.24}$$

$$D_l = -i^{l+1} \frac{W_r\left[\Pi^{(2)}(\eta, r), \Xi(\eta, r)\right]}{W_r\left[\Pi^{(1)}(\eta, r), \Pi^{(2)}(\eta, r)\right]}, \tag{5.25}$$

where the Wronskian determinant $W_x[f(x), g(x)] = f(x) g'(x) - f'(x) g(x)$.

The S-matrix can be obtained by Eqs. (5.24) and (5.25):

$$S_l(\eta) = e^{2i\delta_l} = \frac{D_l}{C_l} = -(-1)^{l+1} \frac{W_r\left[\Pi^{(2)}(\eta, r), \Xi(\eta, r)\right]}{W_r\left[\Pi^{(1)}(\eta, r), \Xi(\eta, r)\right]}. \tag{5.26}$$

The scattering phase shift then reads

$$\delta_l(\eta) = \frac{1}{2} \arg\left(-(-1)^{l+1} \frac{W_r\left[\Pi^{(2)}(\eta, r), \Xi(\eta, r)\right]}{W_r\left[\Pi^{(1)}(\eta, r), \Xi(\eta, r)\right]}\right). \tag{5.27}$$

The scattering phase shift (5.27) can be rewritten as

$$\delta_l(\eta) = W_r\left[\Pi^{(2)}(\eta, r), \Xi(\eta, r)\right] + \frac{l\pi}{2}, \tag{5.28}$$

since $W_r\left[\Pi^{(1)}(\eta, r), \Xi(k, r)\right]$ and $W_r\left[\Pi^{(2)}(\eta, r), \Xi(\eta, r)\right]$ are complex conjugate to each other.

5.4 Asymptotic behavior of scattering wave function

In scattering, we often concern the large-distance asymptotic behavior of the scattering wave function. The asymptotics at $r \to \infty$ of the scattering wave function (5.23) can be achieved in virtue of Eqs. (5.4) and (5.5):

$$\Xi(\eta, r) \overset{r\to\infty}{\sim} \frac{1}{r} \sin\left(\eta r + 2\eta M \ln\frac{r}{M} + M\frac{\mu^2}{\eta} \ln\left(\frac{r}{M} - 1\right) + \delta_l + \chi(\eta, M) - \frac{l\pi}{2} - \eta M\right). \tag{5.29}$$

6 Bound-state eigenvalue: Exact solution

Now we return to the problem of bound states. Based on the solution of scattering obtained above, we present the eigenvalue of the bound state.

In section 4.2, we provide an explicit asymptotic expression for the eigenvalue of bound states. Here we provide an implicit exact expression for the eigenvalue of bound states.

The eigenvalue spectrum of bound states, according to the S-matrix theory, is the singularity of the S-matrix on the positive real axis [47]. Taking $\eta = ik$, we have

$$S_l(k) = -(-1)^{l+1} \frac{W_r\left[\Pi^{(2)}(k,r), \Xi(k,r)\right]}{W_r\left[\Pi^{(1)}(k,r), \Xi(k,r)\right]}. \tag{6.1}$$

The singularity of the S-matrix (6.1) is just the zero of the denominator $W_r\left[\Pi^{(1)}(k,r), \Xi(k,r)\right]$ with $k > 0$, i.e.,

$$W_r\left[\Pi^{(1)}(k,r), \Xi(k,r)\right] = 0, \ k > 0. \tag{6.2}$$

This is an implicit expression of the eigenvalue of bound states. The bound-state eigenvalue $-k^2$ can be solved from Eq. (6.2).

7 Discontinuous jump on horizon: Jump condition

The scattering wave function on the horizon $r = 2M$ has a discontinuous jump. In order to calculate the scattering wave function, we need to first determine the jump of the wave function on the horizon, i.e., the jump condition.

On the horizon, a discontinuous jump of the wave function occurs. In literature, such a discontinuous jump is obtained by an analytic extension [3, 4]. In the following, we will show that by the exact result obtained in the present paper, we can obtain the discontinuous jump through a direct calculation rather than the analytic extension.

7.1 Jump condition: Outgoing wave function

In section 5.2.2, we obtain the exact solutions of the outgoing wave function inside and outside the horizon, Eqs. (5.11) and (5.13), respectively. Comparing the wave functions inside and outside the horizon, we can directly obtain the jump of the wave function on the horizon.

By the outgoing solutions (5.11) and (5.13), the jump of the outgoing wave function on the horizon $r = 2M$ is

$$\left.\frac{\beta^{outer}(u,r)}{\beta^{inner}(u,r)}\right|_{r\to 2M} = \lim_{r\to 2M} \frac{1}{(r-2M)^{-i4\sqrt{\eta^2+\mu^2}M}} = e^{-4M\sqrt{\eta^2+\mu^2}\pi}, \tag{7.1}$$

where $\mathrm{Hc}^{(a)}(\alpha,\beta,\gamma,\delta,\sigma,0) = 1$ [44] is used in the calculation.

It can be directly seen that there exists a jump of the phase of the scattering wave function.

In literature, e.g., Refs. [3, 4], the jump is calculated by the analytic extension.

7.2 Jump condition: Ingoing wave function

Similarly, by the exact solution of the ingoing scattering wave functions inside and outside the horizon obtained in section 5.2.2, we can directly calculate the jump on the horizon of the ingoing wave function.

By the ingoing solutions (5.20) and (5.21), the jump of the ingoing wave function on the horizon $r = 2M$ is

$$\left.\frac{\alpha^{outer}(v,r)}{\alpha^{inner}(v,r)}\right|_{r\to 2M} = \lim_{r\to 2M}(r-2M)^{4i\sqrt{\eta^2+\mu^2}M} = e^{-4M\sqrt{\eta^2+\mu^2}\pi}. \tag{7.2}$$

Comparing Eqs. (7.1) and (7.2) shows that the jump of the outgoing wave and the jump of the ingoing wave are the same.

8 Discontinuous jump on horizon: Comparison with Hawking and Damour-Ruffini methods

In order to calculate the Hawking radiation, one needs to know the discontinuous jump of the wave function on the horizon.

Hawking and Ruffini dealt with this problem from different viewpoints respectively. They, mathematically speaking, did almost the same thing. The method they used is the analytic extension.

In the above, we have obtained the exact wave functions inner and outer the horizon, so we can calculate the jump directly, rather than the analytic extension.

It should be emphasized that the starting point of all the methods is the solution of the outgoing wave of the radial equation (2.2).

In the following, we compare our method with the Hawking treatment and the Damour and Ruffini treatment.

8.1 Hawking method: Brief review

Hawking [3] and Damour and Ruffini [4] both start from the $r \to \infty$ ($r_* \to \infty$) asymptotics of the outgoing wave function

$$\beta^{outer} \overset{r\to\infty}{\sim} \frac{1}{r}e^{-i\omega u} = \frac{1}{r}e^{2i\omega r_*}e^{-i\omega v}, \tag{8.1}$$

where u and v are the outgoing and ingoing Eddington-Finkelstein coordinates, respectively.

Hawking in Ref. [3] uses quantum field theory to deal with the radiation of a black hole.

In this treatment, the outgoing wave function $\Phi_\omega^{outer} = \beta^{outer} Y_{lm}(\theta,\phi)$ is expanded as

$$\Phi_\omega^{outer} = \int d\omega' \left(a_{\omega\omega'} f_{\omega'} + b_{\omega\omega'} \bar{f}_{\omega'}\right) \tag{8.2}$$

with the expansion coefficients

$$a_{\omega\omega'} \sim \frac{1}{2\pi}\left(\frac{\omega'}{\omega}\right)\left(-\omega'\right)^{-i\frac{\omega}{\kappa}-1}, \tag{8.3}$$

$$b_{\omega\omega'} = -i\alpha_{\omega(-\omega')}, \tag{8.4}$$

where f_ω and $\bar{f}_\omega$ are the solutions on past null infinity $f^-(t=-\infty, r=+\infty)$ containing only positive frequencies and only negative frequencies, respectively. The number of particles created by the gravitational field and emitted to infinity then reads [3]

$$\langle 0_-|\, N_\omega \,|0_-\rangle = \int d\omega' \left|b_{\omega\omega'}\right|^2. \tag{8.5}$$

Notice that the subscript of $\alpha_{\omega(-\omega')}$ in Eq. (8.4) is $-\omega'$. In order to calculate the integral in Eq. (8.5), we need to analytic extend ω'. In Eq. (8.3) $\omega' = 0$ is a singular point. Hawking in Ref. [3] performs the analytic extension by anticlockwise round this singularity

$$\omega' \to \omega' e^{-i\pi}. \tag{8.6}$$

By virtue of such an analytic extension, Eq. (8.4) becomes

$$a_{\omega(-\omega')} = ie^{-\frac{\omega}{\kappa}} a_{\omega\omega'}. \tag{8.7}$$

That is to say, $a_{\omega\omega'}$ has a discontinuous jump at $\omega = 0$

$$\frac{a_{\omega(-\omega')}}{a_{\omega\omega'}} = ie^{-\omega/\kappa}. \tag{8.8}$$

Accordingly, Hawking arrives at

$$\langle 0_-| N_\omega |0_-\rangle = \frac{1}{e^{\frac{\omega}{kT}} - 1}, \qquad T = \frac{1}{8\pi Mk}. \tag{8.9}$$

8.2 Damour-Ruffini method: Brief review

Damour and Ruffini, also starting from the asymptotic outgoing wave function (8.1), deferent from Hawking, use scattering method to deal with the radiation of a black hole.

Damour and Ruffini rewrite the asymptotic outgoing wave outside the horizon (8.1) as [4]

$$\beta^{outer} \overset{r\to\infty}{\sim} \frac{1}{r} e^{2i\omega r_*} e^{-i\omega v} = \frac{1}{r} e^{2i\omega r} e^{-i\omega v} \left(\frac{r-2M}{2M}\right)$$

to expose the singularity at the horizon $r = 2M$. In order to analytically extend the outgoing wave to the inner of the horizon, they analytically extend $r - 2M$ as [4]

$$\begin{aligned} r - 2M &\to |r - 2M|\, e^{-i\pi} \\ &= (2M - r)\, e^{-i\pi}. \end{aligned} \tag{8.10}$$

The outgoing wave in the horizon then reads

$$\beta^{inner} = \frac{1}{r} e^{4\pi\omega M} e^{2i\omega r_*} e^{-i\omega r}. \tag{8.11}$$

There is also a discontinuous jump

$$\frac{\beta^{outer}}{\beta^{inner}} = e^{-4\pi\omega M}. \tag{8.12}$$

Then they obtain a relative scattering probability [4]

$$P_\omega = e^{-8\pi\omega M} \tag{8.13}$$

and then obtain the Hawking's result:

$$N_\omega = \frac{1}{e^{8\pi\omega M} - 1} = \frac{1}{e^{\frac{\omega}{kT}} - 1}, \quad T = \frac{1}{8\pi Mk}. \tag{8.14}$$

8.3 Direct calculation through exact solution: Comparison

The starting points of Hawking method and Damour-Ruffini method are both the $r \to 2M$ asymptotic outgoing wave function (8.1), though their treatments are somewhat different.

In this paper, using the outgoing Eddington-Finkelstein coordinate, we obtain the exact solution of the outgoing wave function outside the horizon, Eq. (5.11), and the exact solution of the outgoing wave function inside the horizon, Eq. (5.13). This allows us to calculate the jump exactly.

From these two exact solutions, we can calculate the jump (7.1) straightforwardly without any additional treatment. We then arrive at Hawking and Damour-Ruffini's result

$$\left.\frac{\beta^{outer}}{\beta^{inner}}\right|_{r\to 2M} = e^{-4M\sqrt{\eta^2+\mu^2}\pi} = e^{-4M\omega\pi}. \tag{8.15}$$

The result obtained in the present is based on the exact solution rather than the asymptotic solution. Especially, this approach does not need the analytically extension treatment.

9 Integral equation method: Alternative approach for scattering wave function and phase shift

9.1 Integral equation

In this section, we suggest an alternative approach — an integral equation method — to solve the scattering wave function and the scattering phase shift.

Introducing $u_l(r)$ by

$$\phi_l(r) = \frac{u_l(r)}{r} \tag{9.1}$$

and substituting Eq. (9.1) into Eq. (2.2) with a variable substitution $\rho = r/(2M)$ give an equation of $u_l(\rho)$:

$$\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}u_l(\rho)+\left\{(2M\eta)^2-\left(1-\frac{1}{\rho}\right)\left[\frac{l(l+1)}{\rho^2}+\frac{1}{\rho^3}\right]+\frac{(2M\mu)^2}{\rho}\right\}u_l(\rho)=0. \tag{9.2}$$

By introducing an effective potential

$$V_l^{eff}(\rho) = \left(1-\frac{1}{\rho}\right)\left[\frac{l(l+1)}{\rho^2}+\frac{1}{\rho^3}\right]-\frac{(2M\mu)^2}{\rho}, \tag{9.3}$$

we rewrite Eq. (9.2) as

$$\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}u_l(\rho)+(2M\eta)^2 u_l(\rho) = V_l^{eff}(\rho)\,u_l(\rho). \tag{9.4}$$

9.1.1 Outer horizon

In this section, we solve the equation out of the horizon, i.e., $1 < \rho < \infty$ ($2M < r < \infty$).

In order to solve the equation (9.4) by the Green function method, we first solve Eq. (9.4) with $V_l^{eff}(\rho) = 0$, i.e.,

$$\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}\left(1-\frac{1}{\rho}\right)\frac{d}{d\rho}y_l(\rho) + (2M\eta)^2 y_l(\rho) = 0, \tag{9.5}$$

or,

$$\left(1-\frac{1}{\rho}\right)^2\frac{d^2}{d\rho^2}y_l(\rho) + \left(1-\frac{1}{\rho}\right)\frac{1}{\rho^2}\frac{d}{d\rho}y_l(\rho) + (2M\eta)^2 y_l(\rho) = 0, \tag{9.6}$$

where $y_l(\rho)$ is $u_l(\rho)$ with $V_l^{eff}(\rho) = 0$.

For $1 < \rho < \infty$, Eq. (9.6) has two linearly independent solutions:

$$y_l^{(1)}(\rho) = \sin\left(2M\eta\left[\rho + \ln(\rho-1)\right]\right), \tag{9.7}$$

$$y_l^{(2)}(\rho) = \cos\left(2M\eta\left[\rho + \ln(\rho-1)\right]\right). \tag{9.8}$$

The Green function can be constructed as

$$G(\rho,\rho') = C_1(\rho')\, y_l^{(1)}(\rho) + C_2(\rho')\, y_l^{(2)}(\rho), \quad \rho > \rho', \tag{9.9}$$

$$G(\rho,\rho') = 0, \qquad \rho < \rho', \tag{9.10}$$

in order to satisfy the boundary condition that the Green function must be finite at $\rho = 1$ [48].

Continuity requires that [48]

$$\lim_{\epsilon\to 0^+} G(\rho,\rho')\big|_{\rho=\rho'+\epsilon} = \lim_{\epsilon\to 0^+} G(\rho,\rho')\big|_{\rho=\rho'-\epsilon}, \tag{9.11}$$

$$\lim_{\epsilon\to 0^+}\left[\frac{\partial}{\partial\rho}G(\rho,\rho')\Big|_{\rho=\rho'+\epsilon} - \frac{\partial}{\partial\rho}G(\rho,\rho')\Big|_{\rho=\rho'-\epsilon}\right] = \frac{1}{(1-1/\rho)^2}. \tag{9.12}$$

Then we have

$$C_1(\rho')\sin\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right) + C_2(\rho')\cos\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right) = 0, \tag{9.13}$$

$$\begin{aligned} &2M\eta\left(1+\frac{1}{\rho'-1}\right)C_1(\rho')\cos\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right) \\ &- 2M\eta\left(1+\frac{1}{\rho'-1}\right)C_2(\rho')\sin\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right) = \frac{1}{(1-1/\rho)^2}. \end{aligned} \tag{9.14}$$

Solving $C_1(\rho')$ and $C_2(\rho')$ from Eqs. (9.13) and (9.14) and substituting $C_1(\rho')$, $C_2(\rho')$, and Eqs. (9.7) and (9.8) into Eq. (9.9) give the Green function,

$$\begin{aligned} G(\rho,\rho') =& \frac{\rho'\cos\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right)}{2M\eta(\rho'-1)}\sin\left(2M\eta\left[\rho + \ln(\rho-1)\right]\right) \\ &- \frac{\rho'\sin\left(2M\eta\left[\rho' + \ln(\rho'-1)\right]\right)}{2M\eta(\rho'-1)}\cos\left(2M\eta\left[\rho + \ln(\rho-1)\right]\right), \quad \rho > \rho'. \end{aligned} \tag{9.15}$$

By the Green function (9.15), we can establish an integral equation for $u_l(\rho)$:

$$\begin{aligned} u_l(\rho) &= Ay_l^{(1)}(\rho) + By_l^{(2)}(\rho) + \int_1^\rho G(\rho,\rho') V_l^{eff}(\rho') u_l(\rho') d\rho' \\ &= A\sin(2M\eta[\rho+\ln(\rho-1)]) + B\cos(2M\eta[\rho+\ln(\rho-1)]) \\ &+ \frac{\sin(2M\eta[\rho+\ln(\rho-1)])}{2M\eta}\int_1^\rho \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho') u_l(\rho')\rho' d\rho' \\ &- \frac{\cos(2M\eta[\rho+\ln(\rho-1)])}{2M\eta}\int_1^\rho \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho') u_l(\rho')\rho' d\rho'; \end{aligned} \tag{9.16}$$

or, equivalently,

$$\begin{aligned} u_l(\rho) =& A\sin(2M\eta[\rho+\ln(\rho-1)]) + B\cos(2M\eta[\rho+\ln(\rho-1)]) \\ &- \frac{1}{2M\eta}\int_1^\rho d\rho' \frac{\rho'}{\rho'-1}\sin\left(2M\eta\left[\rho'-\rho+\ln\left(\frac{\rho'-1}{\rho-1}\right)\right]\right) V_l^{eff}(\rho') u_l(\rho'). \end{aligned} \tag{9.17}$$

9.1.2 Wave function near horizon: Outer horizon

On the horizon, the scattering wave function has a discontinuous jump, so we require a jump condition to connect the wave functions inside and outside the horizon. In order to connect the wave functions on the horizon, we first determine the wave function near the horizon at the side of the outer horizon.

With the tortoise coordinate $\rho_* = \int d\rho \frac{1}{1-1/\rho} = \rho + \ln(\rho-1)$ (corresponding to $r_* = r + 2M\ln\left(\frac{r}{2M}-1\right)$) [2], the integral equation (9.17) can be rewritten as

$$u_l(\rho) = A\sin(2M\eta\rho_*) + B\cos(2M\eta\rho_*) - \frac{1}{2M\eta}\int_1^\rho \sin\left(2M\eta\left(\rho'_*-\rho_*\right)\right) V_l^{eff}(\rho') u_l(\rho') d\rho'_*. \tag{9.18}$$

Near the outer horizon, i.e., $\rho \to 1^+$, (corresponding to $r \to 2M$) Eq. (9.18) reduces to

$$u_l(\rho)|_{\rho\to 1^+} = C \lim_{\rho\to 1^+} \sin(2M\eta\rho_* + \phi), \tag{9.19}$$

where $\cos\phi = A/\sqrt{A^2+B^2}$, $\sin\phi = B/\sqrt{A^2+B^2}$, and $C = \sqrt{A^2+B^2}$. The superscript $+$ denotes that ρ tends to the horizon from outside.

Rewrite the wave function (9.19) as

$$u_l(\rho)|_{\rho\to 1^+} = \frac{C}{2i}\lim_{\rho\to 1^+}\left[e^{i(2M\eta\rho_*+\phi)} - e^{-i(2M\eta\rho_*+\phi)}\right], \tag{9.20}$$

which includes two parts:

$$u_l(\rho)|^{out}_{\rho\to 1^+} = \frac{C}{2i}e^{i(2M\eta\rho_*+\phi)}, \quad \text{outgoing wave,} \tag{9.21}$$

$$u_l(\rho)|^{in}_{\rho\to 1^+} = \frac{C}{2i}e^{-i(2M\eta\rho_*+\phi)}, \quad \text{ingoing wave.} \tag{9.22}$$

9.1.3 Inner horizon

In this section, we solve the equation in the horizon, i.e., $0<\rho<1$ ($0<r<2M$).

For $0<\rho<1$, the two linearly independent solutions of Eq. (9.6) are

$$y_l^{(1)}(\rho)=\sin\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right), \tag{9.23}$$

$$y_l^{(2)}(\rho)=\cos\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right). \tag{9.24}$$

The Green function then reads

$$G(\rho,\rho')=0, \qquad \rho>\rho', \tag{9.25}$$

$$G(\rho,\rho')=C_1(\rho')\,y_l^{(1)}(\rho)+C_2(\rho')\,y_l^{(2)}(\rho), \quad \rho<\rho', \tag{9.26}$$

in order to satisfy the boundary condition that the Green function must be finite at $\rho=1$.

Substituting the Green functions (9.25) and (9.26) into the connection condition (9.11) and (9.12) gives

$$C_1(\rho')\sin\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)+C_2(\rho')\cos\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)=0, \tag{9.27}$$

$$\begin{aligned}&2M\eta\left(1+\frac{1}{1-\rho'}\right)C_1(\rho')\cos\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)\\&-2M\eta\left(1+\frac{1}{1-\rho'}\right)C_2(\rho')\sin\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)=-\frac{1}{(1-1/\rho)^2}.\end{aligned} \tag{9.28}$$

Solving $C_1(\rho')$ and $C_2(\rho')$ from Eqs. (9.27) and (9.28) and substituting $C_1(\rho')$, $C_2(\rho')$, and Eqs. (9.23) and (9.24) into Eq. (9.26) give the Green function,

$$\begin{aligned}G(\rho,\rho')=&\frac{\rho'\cos\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)}{2M\eta(1-\rho')}\sin\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)\\&-\frac{\rho'\sin\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)}{2M\eta(1-\rho')}\cos\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right), \quad \rho>\rho'.\end{aligned} \tag{9.29}$$

By the Green function (9.29), we can establish an integral equation for the $u_l(\rho)$ for $0<\rho<1$:

$$\begin{aligned}u_l(\rho)=&\,a y_l^{(1)}(\rho)+b y_l^{(2)}(\rho)+\int_\rho^1 G(\rho,\rho')\,V_l^{eff}(\rho')\,u_l(\rho')\,d\rho'\\=&\,a\sin\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)+b\cos\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)\\&+\frac{\sin\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)}{2M\eta}\int_\rho^1\frac{\cos\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)}{1-\rho'}V_l^{eff}(\rho')\,u_l(\rho')\,\rho'd\rho'\\&-\frac{\cos\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)}{2M\eta}\int_\rho^1\frac{\sin\left(2M\eta\left[\rho'+\ln(1-\rho')\right]\right)}{1-\rho'}V_l^{eff}(\rho')\,u_l(\rho')\,\rho'd\rho',\end{aligned} \tag{9.30}$$

or, equivalently,

$$\begin{aligned}u_l(\rho)=&\,a\sin\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)+b\cos\left(2M\eta\left[\rho+\ln(1-\rho)\right]\right)\\&-\frac{1}{2M\eta}\int_\rho^1 d\rho'\frac{\rho'}{1-\rho'}\sin\left(2M\eta\left[\rho'-\rho+\ln\left(\frac{1-\rho'}{1-\rho}\right)\right]\right)V_l^{eff}(\rho')\,u_l(\rho').\end{aligned} \tag{9.31}$$

9.1.4 Wave function near horizon: Inner horizon

As above, in order to connect the wave functions at the horizon, we first determine the wave function near the horizon at the side of the inner horizon.

With the tortoise coordinate $\rho_* = \rho + \ln(1-\rho)$ [2], the integral equation (9.31) can be rewritten as

$$u_l(\rho) = a\sin(2M\eta\rho_*) + b\cos(2M\eta\rho_*) - \frac{1}{2M\eta}\int_\rho^1 \sin\left(2M\eta\left(\rho'_* - \rho_*\right)\right) V_l^{eff}\left(\rho'\right) u_l\left(\rho'\right) d\rho'_*. \tag{9.32}$$

Near the inner horizon, i.e., $\rho \to 1^-$, Eq. (9.32) reduces to

$$u_l(\rho)|_{\rho\to 1^-} = c\lim_{\rho\to 1^-}\sin(2M\eta\rho_* + \varphi), \tag{9.33}$$

where $\cos\varphi = a/\sqrt{a^2+b^2}$, $\sin\varphi = b/\sqrt{a^2+b^2}$ and $c = \sqrt{a^2+b^2}$. The superscript $+$ denotes that ρ tends to the horizon from inside.

Rewrite the wave function (9.33) as

$$u_l(\rho)|_{\rho\to 1^-} = \frac{c}{2i}\lim_{\rho\to 1^-}\left[e^{i(2M\eta\rho_*+\varphi)} - e^{-i(2M\eta\rho_*+\varphi)}\right], \tag{9.34}$$

which includes two parts:

$$u_l(\rho)|^{out}_{\rho\to 1^-} = \frac{c}{2i}e^{i(2M\eta\rho_*+\varphi)}, \quad \text{outgoing wave}, \tag{9.35}$$

$$u_l(\rho)|^{in}_{\rho\to 1^-} = \frac{c}{2i}e^{-i(2M\eta\rho_*+\varphi)}, \quad \text{ingoing wave}. \tag{9.36}$$

9.2 Connect wave functions by jump condition on horizon

In the above, we obtain the scattering wave functions inside the horizon and outside the horizon, respectively. The wave function has a jump on the horizon. In section 7, we obtain the jump condition. In the following, we will use the jump condition to determine the scattering wave function.

9.2.1 Jump condition

Near the horizon $r = 2M$, a massive field behaves as a massless field, i.e., $\mu \sim 0$, which can be seen directly by observing Eq. (2.2). Therefore, the jump condition (7.1) becomes

$$\left.\frac{\beta^{outer}(v,r)}{\beta^{inner}(v,r)}\right|_{r\to 2M} = \lim_{r\to 2M}(r-2M)^{i4M\sqrt{\eta^2+\mu^2}} = e^{-4M\sqrt{\eta^2+\mu^2}\pi} \sim e^{-4M\eta\pi}. \tag{9.37}$$

9.2.2 Relative scattering amplitude

We have so far obtained the scattering wave functions inside and outside the horizon, Eqs. (9.17) and (9.31). Next, we connect these two wave functions by the connection condition (9.37). By the connection condition, we can determine the undetermined coefficients in the scattering wave function. Concretely, we compare the inner wave function and outer wave function near the horizon, Eqs. (9.21) and (9.35), with the jump condition to determine the coefficients.

The jump condition (9.37) is obtained by use of the outgoing Eddington coordinate. In order to impose the jump condition, we first convert the tortoise coordinate in the wave functions (9.21) and (9.35) into the Eddington coordinate by adding the contribution of the time term in Eqs. (9.21) and (9.35):

$$u_l(\rho,t)|^{out}_{\rho\to 1^+} = \frac{C}{2i}e^{-i\eta t}e^{i(2M\eta\rho_*+\phi)} = \frac{C}{2i}e^{-i\eta v}e^{i(4M\eta\rho_*+\phi)}, \tag{9.38}$$

$$u_l(\rho,t)|^{out}_{\rho\to 1^-} = \frac{c}{2i}e^{-i\eta t}e^{i(2M\eta\rho_*+\varphi)} = \frac{c}{2i}e^{-i\eta v}e^{i(4M\eta\rho_*+\varphi)}. \tag{9.39}$$

To compare with the jump condition, by Eqs. (9.38) and (9.39), we calculate the relative scattering amplitude on the horizon:

$$\begin{aligned}\frac{u_l(\rho)|^{out}_{\rho\to 1^+}}{u_l(\rho)|^{out}_{\rho\to 1^-}} &= \frac{Ce^{i\phi}e^{i4M\eta(\rho+\ln(\rho-1))}}{ce^{i\varphi}e^{i4M\eta(\rho+\ln(1-\rho))}}\\ &= \frac{Ce^{i\phi}}{ce^{i\varphi}}(-1)^{i4M\eta\pi}\\ &= \frac{Ce^{i\phi}}{ce^{i\varphi}}e^{-4M\eta\pi}.\end{aligned} \tag{9.40}$$

Comparing Eq. (9.40) with the jump condition (9.37), we have

$$C = c, \quad \phi = \varphi. \tag{9.41}$$

This gives

$$A = a, \quad B = b. \tag{9.42}$$

The integral equations (9.18) and (9.32) then become

$$\begin{aligned}u_l(\rho) &= A\sin(2M\eta\rho_*) + B\cos(2M\eta\rho_*)\\ &\quad - \frac{1}{2M\eta}\int_1^\rho d\rho'_*\sin\left(2M\eta\left(\rho'_*-\rho_*\right)\right)V_l^{eff}\left(\rho'\right)u_l\left(\rho'\right), \quad \text{inner horizon,}\end{aligned} \tag{9.43}$$

$$\begin{aligned}u_l(\rho) &= A\sin(2M\eta\rho_*) + B\cos(2M\eta\rho_*)\\ &\quad - \frac{1}{2M\eta}\int_\rho^1 d\rho'_*\sin\left(2M\eta\left(\rho'_*-\rho_*\right)\right)V_l^{eff}\left(\rho'\right)u_l\left(\rho'\right), \quad \text{outer horizon.}\end{aligned} \tag{9.44}$$

9.3 Alternative expression of scattering phase shift

In this section, based on the integral equation of the wave function constructed above, we suggest an alternative expression of the scattering phase shift. In this section, we only consider the high energy scattering, i.e., $\mu/\eta \ll 1$.

The expression of the scattering phase shift is obtained by comparing the scattering wave function (5.29) with the radial integral equation given in section 9.

9.3.1 Scattering phase shift

Taking $\rho \to \infty$ asymptotics of Eq. (5.29), for $\mu/\eta \ll 1$, we have

$$\begin{aligned} u_l(\rho) &\overset{\rho\to\infty}{\sim} \sin\left(2M\eta(\rho+\ln(\rho-1)) + \delta_l + \chi(\eta,M) - \frac{l\pi}{2} - \eta M + 2M\eta\ln 2\right) \\ &= \sin(2M\eta[\rho+\ln(\rho-1)])\cos(\delta_l+\Delta(\eta,M)) + \cos(2M\eta[\rho+\ln(\rho-1)])\sin(\delta_l+\Delta(\eta,M)), \end{aligned} \tag{9.45}$$

where $\rho = \frac{r}{2M}$ and $\Delta(\eta,M) = \chi(\eta,M) - \frac{l\pi}{2} - \eta M + 2M\eta\ln 2$ are used.

In order to compare with (9.45), we rewrite Eq. (9.16) as

$$\begin{aligned} u_l(\rho) =& \sin(2M\eta[\rho+\ln(\rho-1)])\left[A + \frac{1}{2M\eta}\int_1^\rho \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'\right] \\ &+ \cos(2M\eta[\rho+\ln(\rho-1)])\left[B - \frac{1}{2M\eta}\int_1^\rho \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'\right]. \end{aligned} \tag{9.46}$$

and, then, take $\rho \to \infty$ asymptotics:

$$u_l(\rho) \overset{\rho\to\infty}{\sim} \alpha(\eta,M)\sin(2M\eta[\rho+\ln(\rho-1)]) + \beta(\eta,M)\cos(2M\eta[\rho+\ln(\rho-1)]) \tag{9.47}$$

$$= C\sin(2M\eta[\rho+\ln(\rho-1)] + \phi), \tag{9.48}$$

where

$$\alpha(\eta,M) = A + \frac{1}{2M\eta}\int_1^\infty \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho', \tag{9.49}$$

$$\beta(\eta,M) = B - \frac{1}{2M\eta}\int_1^\infty \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho', \tag{9.50}$$

$\cos\phi = \alpha(\eta,M)/\sqrt{\alpha^2(\eta,M)+\beta^2(\eta,M)}$, $\sin\phi = \beta(\eta,M)/\sqrt{\alpha^2(\eta,M)+\beta^2(\eta,M)}$, and the normalization constant $C = \sqrt{\alpha^2(\eta,M)+\beta^2(\eta,M)}$.

Comparing Eqs. (9.45) and (9.47), we have

$$\begin{aligned} \tan(\delta_l+\Delta(\eta,M)) &= \frac{\beta(\eta,M)}{\alpha(\eta,M)} \\ &= \frac{B - \frac{1}{2M\eta}\int_1^\infty \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'}{A + \frac{1}{2M\eta}\int_1^\infty \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'}, \end{aligned} \tag{9.51}$$

where

$$\begin{aligned} \phi &= \arctan\frac{\beta(\eta,M)}{\alpha(\eta,M)} \\ &= \frac{B - \frac{1}{2M\eta}\int_1^\infty \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'}{A + \frac{1}{2M\eta}\int_1^\infty \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\,u_l(\rho')\,\rho' d\rho'} \end{aligned} \tag{9.52}$$

is used.

Then we arrive at an expression of the scattering phase shift,

$$\delta_l = \arctan\left(\frac{B - \frac{1}{2M\eta}\int_1^\infty \frac{\sin(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l(\rho')\, \rho' d\rho'}{A + \frac{1}{2M\eta}\int_1^\infty \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l(\rho')\, \rho' d\rho'}\right) - \Delta(\eta, M). \quad (9.53)$$

This is a relation between the scattering phase shift δ_l and the scattering wave function $u_l(\rho)$. To determine the scattering phase shift δ_l, we need to calculate $u_l(\rho)$ and determine the constants A and B.

Now we determine the constants A and B.

The boundary condition requires $u_l(0) = 0$, i.e., the wave function $\Xi(\eta, r)$ must be finite at $r = 0$.

By Eq. (9.31) and the relation $B = b$ given by Eq. (9.42), we have

$$u_l(\rho)|_{\rho=0} = B - \frac{1}{2M\eta}\int_0^1 \frac{\sin(2M\eta[\rho' + \ln(1-\rho')])}{1-\rho'} V_l^{eff}(\rho')\, u_l(\rho')\, \rho' d\rho' = 0. \quad (9.54)$$

Then we have

$$B = \frac{1}{2M\eta}\int_0^1 \frac{\sin(2M\eta[\rho' + \ln(1-\rho')])}{1-\rho'} V_l^{eff}(\rho')\, u_l(\rho')\, \rho' d\rho'. \quad (9.55)$$

Then, with the normalization constant C, the constant A can be determined by Eqs. (9.49) and (9.50).

Next, in order to obtain the scattering phase shift by Eq. (9.53), we need to iteratively solve the wave function $u_l(\rho)$.

9.3.2 Scattering phase shift: zeroth-order and first-order

By iteratively solving the integral equation of the wave function $u_l(\rho)$, Eq. (9.17), we can obtain various orders of $u_l(\rho)$. In this section, we first solve the zeroth-order and first-order scattering phase shifts.

Zeroth-order phase shift $\delta_l^{(0)}$ The zeroth-order contribution of Eq. (9.53) is

$$[\tan(\delta_l + \Delta(\eta, M))]^{(0)} = \frac{B^{(0)}}{A^{(0)}}. \quad (9.56)$$

By Eq. (9.55), we have $B^{(0)} = 0$. Then

$$[\tan(\delta_l + \Delta(\eta, M))]^{(0)} = 0. \quad (9.57)$$

The zeroth-order scattering phase shift reads

$$\delta_l^{(0)} = -\Delta(\eta, M) = -\chi(\eta, M) + \frac{l\pi}{2} + \eta M - 2M\eta\ln 2. \quad (9.58)$$

For small $\frac{1}{M\eta}$, we have [44]

$$\chi(\eta, M) = -2M\eta\ln 2 - \frac{l\pi}{2} - 2M\eta\pi + O\left(\frac{1}{M\eta}\right), \quad (9.59)$$

so

$$\delta_l^{(0)} = l\pi + 2M\eta\pi + \eta M. \quad (9.60)$$

First-order phase shift $\delta_l^{(1)}$ The first-order phase shift $\delta_l^{(1)}$ can be obtained by substituting the zeroth-order scattering wave function $u_l^{(0)}(\rho) = A^{(0)} \sin(2M\eta[\rho + \ln(\rho-1)])$ into Eq. (9.53):

$$\delta_l^{(1)} = -\frac{i}{2}\ln\left(-1 + \frac{4M\eta A^{(1)} + 2\int_1^\infty \frac{\cos(2M\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l^{(0)}(\rho')\, \rho' d\rho'}{2M\eta\left(A^{(1)} - iB^{(1)}\right) + \int_1^\infty \frac{\exp(2iM\eta[\rho'+\ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l^{(0)}(\rho')\, \rho' d\rho'}\right). \tag{9.61}$$

Here by Eq. (9.55) and $C = \sqrt{\alpha^2(\eta, M) + \beta^2(\eta, M)}$, we have

$$B^{(1)} = \frac{1}{2M\eta}\int_0^1 \frac{\sin(2M\eta[\rho' + \ln(\rho'-1)])}{1-\rho'} V_l^{eff}(\rho')\, u_l^{(0)}(\rho')\, \rho' d\rho' \tag{9.62}$$

and

$$\begin{aligned} A^{(1)} = &\sqrt{C^2 - \left[B^{(1)} - \frac{1}{2M\eta}\int_1^\infty \frac{\sin(2M\eta[\rho' + \ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l^{(0)}(\rho')\, \rho' d\rho'\right]^2} \\ &- \frac{1}{2M\eta}\int_1^\infty \frac{\cos(2M\eta[\rho' + \ln(\rho'-1)])}{\rho'-1} V_l^{eff}(\rho')\, u_l^{(0)}(\rho')\, \rho' d\rho'. \end{aligned} \tag{9.63}$$

10 Conclusion

An exact solution for a scalar field in the Schwarzschild spacetime is provided. For bound states, we obtain an exact bound-state wave function and an exact for bound-state eigenvalues. For scattering, we obtain an exact scattering wave function and an exact scattering phase shift.

Moreover, besides the exact solutions, we also provide some approximate solutions. For bound states, we provide an explicit asymptotic expression for bound-state eigenvalues. For scattering, using the integral equation method, we provide the zeroth-order and first-order contributions of scattering phase shift.

In literature, the discontinuous jump on horizon of the wave function is obtained by an analytic extension [3, 4]. In this paper, since we have obtained the exact solution of wave functions inside and outside the horizon, we can calculate this discontinuous jump straightforwardly instead of the analytic extension treatment.

The key step in this paper is to impose the boundary conditions. There are three boundary conditions at $r = 0$, $r = 2M$, and $r \to \infty$. The dynamic equation, however, is a second order differential equation. This is, we have to impose three boundary conditions on one second order differential equation. The approach in the present paper is to first solve the regular solution and the irregular solution —- the solutions satisfying the boundary conditions at $r = 0$, $r = 2M$ and at $r = 2M$, $r \to \infty$, respectively. Then connect the regular solution and the irregular solution together. This treatment can be applied to other kinds of spacetime with not only one singular points.

Scattering by a Schwarzschild spacetime is a long-range potential scattering. In this paper, we present two approaches to deal with this scattering: one is based on the exact solution and one is an integral equation method which provides an approximate solution of

the scattering phase shift. In further researches, we can attempt the heat-kernel method to calculate the scattering phase shift using the approach developed in Refs. [49–55]. Moreover, starting from the exact solution, we can investigate the large-distance asymptotic behavior which is important in the scattering problem [6, 40, 41, 43]. Starting from the exact solution given in the present paper, we can also calculate the corresponding heat kernel and the partition function [56]. Using the heat kernel, we can calculate various quantum field theory quantities, such as one-loop effective actions, vacuum energies, etc. [57, 58].

Acknowledgments

We are very indebted to Dr G. Zeitrauman for his encouragement. This work is supported in part by NSF of China under Grant No. 11575125 and No. 11675119.

References

[1] E. Pike and P. Sabatier, *Scattering: Scattering and Inverse Scattering in Pure and Applied Science*. Scattering: Scattering and Inverse Scattering in Pure and Applied Science. Academic Press, 2002.

[2] V. Frolov and I. Novikov, *Black hole physics: basic concepts and new developments*, vol. 96. Springer Science & Business Media, 2012.

[3] S. W. Hawking, *Particle creation by black holes*, *Communications in mathematical physics* **43** (1975), no. 3 199–220.

[4] T. Damour and R. Ruffini, *Black-hole evaporation in the klein-sauter-heisenberg-euler formalism*, *Physical Review D* **14** (1976), no. 2 332.

[5] J. Futterman, F. Handler, and R. Matzner, *Scattering from Black Holes*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2009.

[6] H. Koyama and A. Tomimatsu, *Asymptotic tails of massive scalar fields in a schwarzschild background*, *Physical Review D* **64** (2001), no. 4 044014.

[7] N. Andersson and K.-E. Thylwe, *Complex angular momentum approach to black-hole scattering*, *Classical and Quantum Gravity* **11** (1994), no. 12 2991.

[8] C. F. Macedo and L. C. Crispino, *Absorption of planar massless scalar waves by bardeen regular black holes*, *Physical Review D* **90** (2014), no. 6 064001.

[9] D. Batic, N. Kelkar, and M. Nowakowski, *On born approximation in black hole scattering*, *The European Physical Journal C* **71** (2012), no. 12 1–8.

[10] N. Andersson, *Scattering of massless scalar waves by a schwarzschild black hole: A phase-integral study*, *Physical Review D* **52** (1995), no. 4 1808.

[11] J. G. Rosa and S. R. Dolan, *Massive vector fields on the schwarzschild spacetime: quasinormal modes and bound states*, *Physical Review D* **85** (2012), no. 4 044043.

[12] W. M. Jin, *Scattering of massive dirac fields on the schwarzschild black hole spacetime*, *Classical and Quantum Gravity* **15** (1998), no. 10 3163.

[13] M. Y. Kuchiev and V. Flambaum, *Scattering of scalar particles by a black hole*, *Physical Review D* **70** (2004), no. 4 044022.

[14] D. Batic, N. Kelkar, and M. Nowakowski, *Orbiting phenomena in black hole scattering*, *Physical Review D* **86** (2012), no. 10 104060.

[15] S. Dolan, C. Doran, and A. Lasenby, *Fermion scattering by a schwarzschild black hole*, *Physical Review D* **74** (2006), no. 6 064005.

[16] H. Cho and Y. Lin, *Wkb analysis of the scattering of massive dirac fields in schwarzschild black-hole spacetimes*, *Classical and Quantum Gravity* **22** (2005), no. 5 775.

[17] C. L. Benone, E. S. de Oliveira, S. R. Dolan, and L. C. Crispino, *Absorption of a massive scalar field by a charged black hole*, *Physical Review D* **89** (2014), no. 10 104053.

[18] L. C. Crispino, S. R. Dolan, and E. S. Oliveira, *Scattering of massless scalar waves by reissner-nordström black holes*, *Physical Review D* **79** (2009), no. 6 064022.

[19] G. Pei and C. Bambi, *Scattering of particles by deformed non-rotating black holes*, *The European Physical Journal C* **75** (2015), no. 11 1–7.

[20] K. Glampedakis and N. Andersson, *Scattering of scalar waves by rotating black holes*, *Classical and Quantum Gravity* **18** (2001), no. 10 1939.

[21] C. Doran and A. Lasenby, *Perturbation theory calculation of the black hole elastic scattering cross section*, *Physical Review D* **66** (2002), no. 2 024006.

[22] R. Brito, V. Cardoso, and P. Pani, *Massive spin-2 fields on black hole spacetimes: Instability of the schwarzschild and kerr solutions and bounds on the graviton mass*, *Physical Review D* **88** (2013), no. 2 023514.

[23] D. Ahn, Y. Moon, R. Mann, and I. Fuentes-Schuller, *The black hole final state for the dirac fields in schwarzschild spacetime*, *Journal of High Energy Physics* **2008** (2008), no. 06 062.

[24] V. Cardoso and J. P. Lemos, *Quasinormal modes of schwarzschild–anti-de sitter black holes: Electromagnetic and gravitational perturbations*, *Physical Review D* **64** (2001), no. 8 084017.

[25] M. Giammatteo and J. Jing, *Dirac quasinormal frequencies in schwarzschild-ads space-time*, *Physical Review D* **71** (2005), no. 2 024007.

[26] H. Okawa, K.-i. Nakao, and M. Shibata, *Is super-planckian physics visible? scattering of black holes in 5 dimensions*, *Physical Review D* **83** (2011), no. 12 121501.

[27] R. Konoplya and E. Abdalla, *Scalar field perturbations of the schwarzschild black hole in the gödel universe*, *Physical Review D* **71** (2005), no. 8 084015.

[28] T. Damour and R. Ruffini, *Quantum electrodynamical effects in kerr-newmann geometries*, *Physical Review Letters* **35** (1975), no. 7 463.

[29] R. Zhao, L.-C. Zhang, and H.-F. Li, *Hawking radiation of a reissner–nordström–de sitter black hole*, *General Relativity and Gravitation* **42** (2010), no. 4 975–983.

[30] X. He and W. Liu, *Modified hawking radiation in a btz black hole using damour–ruffini method*, *Physics Letters B* **653** (2007), no. 2 330–334.

[31] S. Zhou and W. Liu, *Hawking radiation of charged dirac particles from a kerr-newman black hole*, *Physical Review D* **77** (2008), no. 10 104021.

[32] S. Sannan, *Heuristic derivation of the probability distributions of particles emitted by a black hole*, *General Relativity and Gravitation* **20** (1988), no. 3 239–246.

[33] R. Li, J.-R. Ren, and S.-W. Wei, *Hawking radiation of dirac particles via tunneling from the kerr black hole*, *Classical and Quantum Gravity* **25** (2008), no. 12 125016.

[34] R. Li and J.-R. Ren, *Dirac particles tunneling from btz black hole*, *Physics Letters B* **661** (2008), no. 5 370–372.

[35] L.-C. Zhang, H.-F. Li, and R. Zhao, *Hawking radiation from a rotating acoustic black hole*, *Physics Letters B* **698** (2011), no. 5 438–442.

[36] R. Balbinot, A. Fabbri, P. Nicolini, V. Frolov, P. Sutton, and A. Zelnikov, *Vacuum polarization in the schwarzschild spacetime and dimensional reduction*, *Physical Review D* **63** (2001), no. 8 084029.

[37] P. Fiziev and D. Staicova, *Application of the confluent heun functions for finding the quasinormal modes of nonrotating black holes*, *Physical Review D* **84** (2011), no. 12 127502.

[38] H. Vieira and V. Bezerra, *Confluent heun functions and the physics of black holes: Resonant frequencies, hawking radiation and scattering of scalar waves*, *Annals of Physics* **373** (2016) 28–42.

[39] H. Vieira, V. Bezerra, and C. Muniz, *Exact solutions of the klein–gordon equation in the kerr–newman background and hawking radiation*, *Annals of Physics* **350** (2014) 14–28.

[40] T. Liu, W.-D. Li, and W.-S. Dai, *Scattering theory without large-distance asymptotics*, *Journal of High Energy Physics* **2014** (2014), no. 6 1–12.

[41] W.-D. Li and W.-S. Dai, *Scattering theory without large-distance asymptotics in arbitrary dimensions*, *Journal of Physics A: Mathematical and Theoretical* **49** (2016), no. 46 465202.

[42] W.-D. Li and W.-S. Dai, *Exact solution of inverse-square-root potential* $v(r) = -\frac{\alpha}{\sqrt{r}}$, *Annals of Physics* **373** (2016) 207 – 215.

[43] S. Hod, *Scattering by a long-range potential*, *Journal of High Energy Physics* **2013** (2013), no. 9 1–11.

[44] A. Ronveaux and F. M. Arscott, *Heun's differential equations*. Oxford University Press, 1995.

[45] H. Friedrich, *Scattering Theory*. Springer Berlin Heidelberg, 2015.

[46] S. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*. Addison Wesley, 2004.

[47] C. J. Joachain, *Quantum collision theory*. North-Holland Publishing Company, Amsterdam, 1975.

[48] G. Arfken, H. Weber, and F. Harris, *Mathematical Methods for Physicists: A Comprehensive Guide*. Elsevier, 2012.

[49] N. Graham, M. Quandt, and H. Weigel, *Spectral methods in quantum field theory*, vol. 777. Springer, 2009.

[50] A. Barvinsky and G. Vilkovisky, *Beyond the schwinger-dewitt technique: Converting loops into trees and in-in currents*, *Nuclear Physics B* **282** (1987) 163–188.

[51] A. Barvinsky and G. Vilkovisky, *Covariant perturbation theory (ii). second order in the curvature. general algorithms*, *Nuclear Physics B* **333** (1990), no. 2 471–511.

[52] A. Barvinsky and G. Vilkovisky, *Covariant perturbation theory (iii). spectral representations of the third-order form factors*, *Nuclear Physics B* **333** (1990), no. 2 512–524.

[53] V. Mukhanov and S. Winitzki, *Introduction to quantum effects in gravity*. Cambridge University Press, 2007.

[54] H. Pang, W.-S. Dai, and M. Xie, *Relation between heat kernel method and scattering spectral method*, *The European Physical Journal C* **72** (2012), no. 5 1–13.

[55] W.-D. Li and W.-S. Dai, *Heat-kernel approach for scattering*, *The European Physical Journal C* **75** (2015), no. 6.

[56] D. V. Vassilevich, *Heat kernel expansion: user's manual*, *Physics reports* **388** (2003), no. 5 279–360.

[57] W.-S. Dai and M. Xie, *The number of eigenstates: counting function and heat kernel*, *Journal of High Energy Physics* **2009** (2009), no. 02 033.

[58] W.-S. Dai and M. Xie, *An approach for the calculation of one-loop effective actions, vacuum energies, and spectral counting functions*, *Journal of High Energy Physics* **2010** (2010), no. 6 1–29.